
COMPARISON OF LUCAS AND THE NATIONAL SOIL MONITORING PROGRAMMES IN SWEDEN



Johanna Wetterlind, johanna.Wetterlind@slu.se SLU, Department of Soil and Environment





Background

A review of existing soil monitoring systems that was done within EJP SOIL (WP6), demonstrated that soil monitoring systems across EU countries (including LUCAS) are quite different (e.g. sampling strategies and methods, analysis of soil samples...)

• Deliverable 6.3, https://ejpsoil.eu/fileadmin/projects/ejpsoil/WP6/EJP_SOIL_Deliverable_6.3_Dec_2021_final.pdf

Two exercises were developed (lead by INRAE)

- The comparison of LUCAS dataset with national data, country by country
 - Present some of the Swedish results
 - Paper soon to be submitted
- A double sampling exercise to develop transfer functions, in collaboration with JRC
 - Analytical procedures: LUCAS samples analysed at "LUCAS lab" and national lab 17 countries
 - Sampling and analytical procedures: sampling according to both protocols 6 countries




Sample density and distribution




LUCAS 2015


National ~2015

LUCAS 2022









Total: 1 847

Cropland: 154 (8%)

Forest: 1 496 (81%)

6 327

2 030 (32%)

4 297 (68%)

2 845


759 (27%)

1 463 (51%)




Soil type and land use





1986, Troedsson, T.; Wiberg, M., KSLA, ESDAC



Soil samples in agricultural land



Problem with classification of grasslands



Share of total area by type and land cover (%), 2018

	Total area (km2)	Woodland and shrubland	Cropland	Grassland	Water areas and wetland; bareland	Artificial
EU	4 125 107	46.8	24.2	17.4	7.3	4.2
Finland	338 411	69.6	5.3	5.7	17.6	1.7
Sweden	447 424	68.5	4.0	5.5	20.1	1.8

Source: Eurostat (online data code: lan_lcv_ovw)

eurostat

Fodder crops: 0.9 %



Swedish national statistics (Land Use in Sweden 2020, Statistics Sweden (SCB))

Agricultural land: 7 %

- 6 % cropland
- 1 % grassland (pastures)
- 45 % of the cropland was ley (often grass clover mix), about 2.7 % of the total land use

Ley: temporal grassland (often 1-3 year grass, clover or grass-clover mix) in the crop rotation, on cropland.





Problem with classification of grasslands



It looks like part of the land classified as grassland in LUCAS is ley in a crop rotation on crop land.

Why is that a problem?

- Statistics related to important land use changes becomes unreliable.
 - Grassland to cropland
 - Impact on scenario modelling
- Part of the croplands are not included in the cropland statistics
 - Biased excluding soils that benefits from having ley in the rotation
 - and often organic fertiliser inputs potentially with god soil health
- Difficult to compare European and national statistics






Some soil properties




Soil texture

LUCAS (2015)
(cropland + grassland without trees)



Swedish Soil and crop monitoring programme (cropland)







Some soil properties



Organic carbon









Challenges for harmonisation



Main issue is to be able to continue to use the existing inventories and time series, while at the same time provide good data for the European scale.

- Soil depth
 - LUCAS (since 2022, 0-30) Soils and crop inventory (0-20), Forest Soil inventory (several depth)
- Sampling protocol
 - Especially for the forest soils
- Different analyses methods
 - e.g. plant available nutrients
 - Also want be relevant for national agriculture advise

Important not to loos the national relevance to be able to fit into a European framework.



Possibilities for the national soil monitoring



Increased interest and visibility of the national inventories!

Possibilities to include new analyses that are not part of the current inventories.

- e.g. Bulk density in agricultural soils?
- e.g. P-Olsen in parallel with P-AL?

Develop new inventories for better national information on soil erosion and water related properties?

